

# North Star School District Concussion Guide

- Parents Guide To Concussion In Sports Pages 2-7
- Suggested Guidelines for Management of Concussion Page 8
- Suggested Guidelines for Management of Concussions in Sports Page 9-13  
**National Federation of State High School Associations (NFHS)**  
**Sports Medicine Advisory Committee (SMAC)**
- Suggested Guidelines for Management of Concussion Page 14-15  
**(Brochure)**



## **A Parent's Guide to Concussion in Sports**

### **What is a concussion?**

- A concussion is a brain injury which results in a temporary disruption of normal brain function. A concussion occurs when the brain is violently rocked back and forth or twisted inside the skull as a result of a blow to the head or body. An athlete does not have to lose consciousness (“knocked-out”) to suffer a concussion.

### **Concussion Facts**

- It is estimated that over 140,000 high school athletes across the United States suffer a concussion each year. (Data from NFHS Injury Surveillance System)
- Concussions occur most frequently in football, but girl's lacrosse, girl's soccer, boy's lacrosse, wrestling and girl's basketball follow closely behind. All athletes are at risk.
- A concussion is a traumatic injury to the brain.
- Concussion symptoms may last from a few days to several months.
- Concussions can cause symptoms which interfere with school, work, and social life.
- An athlete should not return to sports while still having symptoms from a concussion as they are at risk for prolonging symptoms and further injury.
- A concussion may cause multiple symptoms. Many symptoms appear immediately after the injury, while others may develop over the next several days or weeks. The symptoms may be subtle and are often difficult to fully recognize.

## What are the signs and symptoms of a concussion?

### SIGNS OBSERVED BY PARENTS, FRIENDS, TEACHERS OR COACHES

Appears dazed or stunned

Is confused about what to do

Forgets plays

Is unsure of game, score, or opponent

Moves clumsily

Answers questions slowly

Loses consciousness

Shows behavior or personality changes

Can't recall events prior to hit

Can't recall events after hit

### SYMPTOMS REPORTED BY ATHLETE

Headache

Nausea

Balance problems or dizziness

Double or fuzzy vision

Sensitivity to light or noise

Feeling sluggish

Feeling foggy or groggy

Concentration or memory problems

Confusion

## What should I do if I think my child has had a concussion?

If an athlete is suspected of having a concussion, he or she must be immediately removed from play, be it a game or practice. Continuing to participate in physical activity after a concussion can lead to worsening concussion symptoms, increased risk for further injury, and even death. Parents and coaches are not expected to be able to “diagnose” a concussion, as that is the job of a medical professional. However, you must be aware of the signs and symptoms of a concussion and if you are suspicious, then your child must stop playing:

### When in doubt, sit them out!

All athletes who sustain a concussion need to be evaluated by a health care professional who is familiar with sports concussions. You should call your child's physician and explain what has happened and follow your physician's instructions. If your child is vomiting, has a severe headache, is having difficulty staying awake or answering simple questions he or she should be taken to the emergency department immediately.

## **When can an athlete return to play following a concussion?**

After suffering a concussion, **no athlete should return to play or practice on that same day.** Previously, athletes were allowed to return to play if their symptoms resolved within 15 minutes of the injury. Studies have shown us that the young brain does not recover quickly enough for an athlete to return to activity in such a short time.

Concerns over athletes returning to play too quickly have led state lawmakers in both Oregon and Washington to pass laws stating that **no player shall return to play following a concussion on that same day and the athlete must be cleared by an appropriate health-care professional before he or she are allowed to return to play in games or practices.** The laws also mandate that coaches receive education on recognizing the signs and symptoms of concussion.

Once an athlete no longer has symptoms of a concussion and is cleared to return to play by health care professional knowledgeable in the care of sports concussions he or she should proceed with activity in a step-wise fashion to allow the brain to re-adjust to exertion. On average the athlete will complete a new step each day. The return to play schedule should proceed as below following medical clearance:

*Step 1:* Light exercise, including walking or riding an exercise bike. No weight-lifting.

*Step 2:* Running in the gym or on the field. No helmet or other equipment.

*Step 3:* Non-contact training drills in full equipment. Weight-training can begin.

*Step 4:* Full contact practice or training.

*Step 5:* Game play.

**If symptoms occur at any step, the athlete should cease activity and be re-evaluated by their health care provider.**

## **How can a concussion affect schoolwork?**

Following a concussion, many athletes will have difficulty in school. These problems may last from days to months and often involve difficulties with short and long-term memory, concentration, and organization.

In many cases it is best to lessen the athlete's class load early on after the injury. This may include staying home from school for a few days, followed by a lightened schedule for a few days, or perhaps a longer period of time, if needed. Decreasing the stress on the brain early on after a concussion may lessen symptoms and shorten the recovery time.

## **What can I do?**

- Both you and your child should learn to recognize the “Signs and Symptoms” of concussion as listed above.
- Teach your child to tell the coaching staff if he or she experiences such symptoms.
- Emphasize to administrators, coaches, teachers, and other parents your concerns and expectations about concussion and safe play.
- Teach your child to tell the coaching staff if he or she suspects that a teammate has a concussion.
- Monitor sports equipment for safety, fit, and maintenance.
- Ask teachers to monitor any decrease in grades or changes in behavior that could indicate concussion.
- Report concussions that occurred during the school year to appropriate school staff. This will help in monitoring injured athletes as they move to the next season’s sports.

## **Other Frequently Asked Questions**

### **Why is it so important that an athlete not return to play until they have completely recovered from a concussion?**

Athletes who are not fully recovered from an initial concussion are significantly vulnerable for recurrent, cumulative, and even catastrophic consequences of a second concussive injury. Such difficulties are prevented if the athlete is allowed time to recover from the concussion and return to play decisions are carefully made. No athlete should return-to-sport or other at-risk participation when symptoms of concussion are present and recovery is ongoing.

### **Is a “CAT scan” or MRI needed to diagnose a concussion?**

Diagnostic testing, which includes CT (“CAT”) and MRI scans, are rarely needed following a concussion. While these are helpful in identifying life-threatening brain injuries (e.g. skull fracture, bleeding, swelling), they are not normally utilized, even by athletes who have sustained severe concussions. A concussion is diagnosed based upon the athlete’s story of the injury and the health care provider’s physical examination.

### **What is the best treatment to help my child recover more quickly from a concussion?**

The best treatment for a concussion is rest. There are no medications that can speed the recovery from a concussion. Exposure to loud noises, bright lights, computers, video games, television and phones (including text messaging) all may worsen the symptoms of a concussion. You should allow your child to rest as much as possible in the days following a concussion. As the symptoms

lessen, you can allow increased use of computers, phone, video games, etc., but the access must be lessened if symptoms worsen.

### **How long do the symptoms of a concussion usually last?**

The symptoms of a concussion will usually go away within one week of the initial injury. You should anticipate that your child will likely be out of sports for about two weeks following a concussion. However, in some cases symptoms may last for several weeks, or even months. Symptoms such as headache, memory problems, poor concentration, and mood changes can interfere with school, work, and social interactions. The potential for such long-term symptoms indicates the need for careful management of all concussions.

### **How many concussions can an athlete have before he or she should stop playing sports?**

There is no “magic number” of concussions that determine when an athlete should give up playing contact or collision sports. The circumstances surrounding each individual injury, such as how the injury happened and length of symptoms following the concussion, are very important and must be considered when assessing an athlete’s risk for further and potentially more serious concussions. The decision to “retire” from sports is a decision best reached following a complete evaluation by your child’s primary care provider and consultation with a physician or neuropsychologist who specializes in treating sports concussion.

### **I’ve read recently that concussions may cause long-term brain damage in professional football players. Is this a risk for high school athletes who have had a concussion?**

The issue of “chronic encephalopathy” in several former NFL players has received a great deal of media attention lately. Very little is known about what may be causing dramatic abnormalities in the brains of these unfortunate retired football players. At this time we have very little knowledge of the long-term effects of concussions which happen during high school athletics.

In the cases of the retired NFL players, it appears that most had long careers in the NFL after playing in high school and college. In most cases, they played football for over 20 years and suffered multiple concussions in addition to hundreds of other blows to their heads. Alcohol and steroid use may also be contributing factors in some cases. Obviously, the average high school athlete does not come close to suffering the total number or shear force of head trauma seen by professional football players. However, the fact that we know very little about the long-term effects of concussions in young athletes is further reason to very carefully manage each concussion.

Some of this information has been adapted from the CDC's "Heads Up: Concussion in High School Sports" materials by the NFHS's Sports Medicine Advisory Committee. Please go to [www.cdc.gov/ncipc/tbi/Coaches\\_Tool\\_Kit.htm](http://www.cdc.gov/ncipc/tbi/Coaches_Tool_Kit.htm) for more information.

If you have any further questions regarding concussions in high school athletes or want to know how to find a concussion specialist in your area please contact Michael C. Koester, MD, ATC and Chair of the NFHS Sports Medicine Advisory Committee at [michael.koester@slocumcenter.com](mailto:michael.koester@slocumcenter.com).

**April 2010**



# Suggested Guidelines for Management of Concussion

A concussion is a traumatic brain injury that interferes with normal brain function. An athlete does not have to lose consciousness (be “knocked out”) to have suffered a concussion.

## **Common Symptoms of Concussion Include:**

- headache
- fogginess
- difficulty concentrating
- easily confused
- slowed thought processes
- difficulty with memory
- nausea
- lack of energy, tiredness
- dizziness, poor balance
- blurred vision
- sensitive to light and sounds
- mood changes- irritable, anxious, or tearful

## **Suggested Concussion Management:**

1. No athlete should return to play (RTP) or practice on the same day of a concussion.
2. Any athlete suspected of having a concussion should be evaluated by an appropriate health-care professional that day.
3. Any athlete with a concussion should be medically cleared by an appropriate health-care professional prior to resuming participation in any practice or competition.
4. After medical clearance, RTP should follow a step-wise protocol with provisions for delayed RTP based upon return of any signs or symptoms.

For further details please see the “NFHS Suggested Guidelines for Management of Concussion” at [www.nfhs.org](http://www.nfhs.org).



# SUGGESTED GUIDELINES FOR MANAGEMENT OF CONCUSSION IN SPORTS

## National Federation of State High School Associations (NFHS) Sports Medicine Advisory Committee (SMAC)

### Introduction

A concussion is type of traumatic brain injury that interferes with normal function of the brain. It occurs when the brain is rocked back and forth or twisted inside the skull as a result of a blow to the head or body. What may appear to be only a mild jolt or blow to the head or body can result in a concussion.

The understanding of sports-related concussion has evolved dramatically in recent years. We now know that young athletes are particularly vulnerable to the effects of a concussion. Once considered little more than a “ding” on the head, it is now understood that a concussion has the potential to result in short or long-term changes in brain function, or in some cases, death.

### What is a concussion?

You’ve probably heard the terms “ding” and “bell-ringer.” These terms were once used to refer to minor head injuries and thought to be a normal part of sports. There is no such thing as a minor brain injury. Any suspected concussion must be taken seriously. A concussion is caused by a bump, blow, or jolt to the head or body. Basically, any force that is transmitted to the head causes the brain to literally bounce around or twist within the skull, potentially resulting in a concussion.

**It used to be believed that a player had to lose consciousness or be “knocked-out” to have a concussion. This is not true, as the vast majority of concussions do not involve a loss of consciousness. In fact, less than 10% of players actually lose consciousness with a concussion.**

What exactly happens to the brain during a concussion is not entirely understood. It appears to be a very complex injury affecting both the structure and function of the brain. The sudden movement of the brain causes stretching and tearing of brain cells, damaging the cells and creating chemical changes in the brain. Once this injury occurs, the brain is vulnerable to further injury and very sensitive to any increased stress until it fully recovers.

Common sports injuries such as torn ligaments and broken bones are structural injuries that can be seen on MRIs or x-rays, or detected during an examination. A concussion, however, is primarily an injury that interferes with how the brain works. While there is damage to brain cells, the damage is at a microscopic level and cannot be seen on MRI or CT scans. Therefore, the brain looks normal on these tests, even though it has been seriously injured.

## Recognition and Management

If an athlete exhibits any signs, symptoms, or behaviors that make you suspicious that he or she may have had a concussion, that athlete must be removed from all physical activity, including sports and recreation. Continuing to participate in physical activity after a concussion can lead to worsening concussion symptoms, increased risk for further injury, and even death.

<b>SYMPTOMS REPORTED BY ATHLETE</b>
Headache
Nausea
Balance problems or dizziness
Double or fuzzy vision
Sensitivity to light or noise
Feeling sluggish
Feeling foggy or groggy
Concentration or memory problems
Confusion

Parents and coaches are not expected to be able to “diagnose” a concussion. That is the role of an appropriate health-care professional. However, you must be aware of the signs, symptoms and behaviors of a possible concussion, and if you suspect that an athlete may have a concussion, then he or she must be immediately removed from all physical activity.

<b>SIGNS OBSERVED BY PARENTS, FRIENDS, TEACHERS OR COACHES</b>
Appears dazed or stunned
Is confused about what to do
Forgets plays
Is unsure of game, score, or opponent
Moves clumsily
Answers questions slowly
Loses consciousness
Shows behavior or personality changes
Can't recall events prior to hit
Can't recall events after hit

**When in doubt, sit them out!**

When you suspect that a player has a concussion, follow the “Heads Up” 4-step Action Plan.

1. Remove the athlete from play.
2. Ensure that the athlete is evaluated by an appropriate health-care professional.
3. Inform the athlete’s parents or guardians about the possible concussion and give them information on concussion.
4. Keep the athlete out of play the day of the injury and until an appropriate health-care professional says he or she is symptom-free and gives the okay to return to activity.

The signs, symptoms, and behaviors of a concussion are not always apparent immediately after a bump, blow, or jolt to the head or body and may develop over a few hours. An athlete should be observed following a suspected concussion and should never be left alone.

Athletes must know that they should never try to “tough out” a suspected concussion. Teammates, parents and coaches should never encourage an athlete to “play through” the symptoms of a concussion. In addition, there should never be an attribution of bravery to athletes who do play despite having concussion signs or symptoms. The risks of such behavior must be emphasized to all members of the team, as well as coaches and parents.

If an athlete returns to activity before being fully healed from an initial concussion, the athlete is at risk for a repeat concussion. A repeat concussion that occurs before the brain has a chance to recover from the first can slow recovery or increase the chance for long-term problems. In rare cases, a repeat concussion can result in severe swelling and bleeding in the brain that can be fatal.

### **Cognitive Rest**

A concussion can interfere with school, work, sleep and social interactions. Many athletes who have a concussion will have difficulty in school with short- and long-term memory, concentration and organization. These problems typically last no longer than a week or two, but for some these difficulties may last for months. It is best to lessen the student’s class load early on after the injury. Most students with concussion recover fully. However, returning to sports and other regular activities too quickly can prolong the recovery.

The first step in recovering from a concussion is rest. Rest is essential to help the brain heal. Students with a concussion need rest from physical and mental activities that require concentration and attention as these activities may worsen symptoms and delay recovery. Exposure to loud noises, bright lights, computers, video games, television and phones (including texting) all may worsen the symptoms of concussion. As the symptoms lessen, increased use of computers, phone, video games, etc., may be allowed.

### **Return to Play**

After suffering a concussion, **no athlete should return to play or practice on that same day.** Previously, athletes were allowed to return to play if their symptoms resolved within 15 minutes of the injury. Newer studies have shown us that the young brain does not recover quickly enough for an athlete to return to activity in such a short time.

**An athlete should never be allowed to resume physical activity following a concussion until he or she is symptom free and given the approval to resume physical activity by an appropriate health-care professional.**

Once an athlete no longer has signs, symptoms, or behaviors of a concussion **and is cleared to return to activity by a health-care professional**, he or she should proceed in a step-wise fashion to allow the brain to re-adjust to exercise. In most cases, the athlete will progress one step each day. The return to activity program schedule **may** proceed as below **following medical clearance**:

### **Progressive Physical Activity Program**

- Step 1:* Light aerobic exercise- 5 to 10 minutes on an exercise bike or light jog; no weight lifting, resistance training, or any other exercises.
- Step 2:* Moderate aerobic exercise- 15 to 20 minutes of running at moderate intensity in the gym or on the field without a helmet or other equipment.
- Step 3:* Non-contact training drills in full uniform. May begin weight lifting, resistance training, and other exercises.
- Step 4:* Full contact practice or training.
- Step 5:* Full game play.

**If symptoms of a concussion re-occur, or if concussion signs and/or behaviors are observed at any time during the return to activity program, the athlete must discontinue all activity and be re-evaluated by their health care provider.**

### **Concussion in the Classroom**

Following a concussion, many athletes will have difficulty in school. These problems may last from days to months and often involve difficulties with short- and long-term memory, concentration, and organization. In many cases, it is best to lessen the student's class load early on after the injury. This may include staying home from school for a few days, followed by a lightened schedule for a few days, or longer, if necessary. Decreasing the stress on the brain early on after a concussion may lessen symptoms and shorten the recovery time.

### **What to do in an Emergency**

Although rare, there are some situations where you will need to call 911 and activate the Emergency Medical System (EMS). The following circumstances are medical emergencies:

1. Any time an athlete has a loss of consciousness of any duration. While loss of consciousness is not required for a concussion to occur, it may indicate more serious brain injury.
2. If an athlete exhibits any of the following: decreasing level of consciousness, looks very drowsy or cannot be awakened, if there is difficulty getting his or her attention, irregularity in breathing, severe or worsening headaches, persistent vomiting, or any seizures.

### **Suggested Concussion Management**

1. No athlete should return to play (RTP) or practice on the same day of a concussion.
2. Any athlete suspected of having a concussion should be evaluated by an appropriate health-care professional that day.
3. Any athlete with a concussion should be medically cleared by an appropriate health-care professional prior to resuming participation in any practice or competition.
4. After medical clearance, RTP should follow a step-wise protocol with provisions for delayed RTP based upon return of any signs or symptoms.

## **References**

Guskiewicz KM, et al. National Athletic Trainers' Association position statement: management of sport-related concussion. Journal of Athletic Training 2004; 39:280-297.

McCrory P, et al. Consensus statement on concussion in sport: the 3rd International Conference on Concussion in Sport held in Zurich, November 2008. Journal of Athletic Training 2009; 44:434-48.

## **Additional Resources**

Heads Up: Concussion in High School Sports

[http://www.cdc.gov/concussion/headsup/high\\_school.html](http://www.cdc.gov/concussion/headsup/high_school.html)

Concussion in Sports- What you need to know.

<http://www.nfhslearn.com/electiveDetail.aspx?courseID=15000>

NFHS Sports Medicine Handbook, 4<sup>th</sup> Ed, 2011.

**Revised January 2011**

## SIGNS AND SYMPTOMS OF CONCUSSION

Concussions can appear in many different ways. Listed below are some of the signs and symptoms frequently associated with concussions. Most signs, symptoms and abnormalities after a concussion fall into the four categories listed below. A coach, parent or other person who knows the athlete well can often detect these problems by observing the athlete and/or by asking a few relevant questions of the athlete, official or a teammate who was on the field or court at the time of the concussion. Below are some suggested observations and questions a non-medical individual can use to help determine whether an athlete has suffered a concussion and how urgently he or she should be sent for appropriate medical care.

### 1. PROBLEMS IN BRAIN FUNCTION:

- a. Confused state – dazed look, vacant stare or confusion about what happened or is happening.
- b. Memory problems – can't remember assignment on play, opponent, score of game, or period of the game; can't remember how or with whom he or she traveled to the game, what he or she was wearing, what was eaten for breakfast, etc.
- c. Symptoms reported by athlete – Headache, nausea or vomiting; blurred or double vision; oversensitivity to sound, light or touch; ringing in ears; feeling foggy or groggy; dizziness.
- d. Lack of sustained attention – difficulty sustaining focus adequately to complete a task, a coherent thought or a conversation.

**2. SPEED OF BRAIN FUNCTION:** Slow response to questions, slow slurred speech, incoherent speech, slow body movements and slow reaction time.

**3. UNUSUAL BEHAVIORS:** Behaving in a combative, aggressive or very silly manner; atypical behavior for the individual; repeatedly asking the same question over and over; restless and irritable behavior with constant motion and attempts to return to play; reactions that seem out of proportion and inappropriate; and having trouble resting or "finding a comfortable position."

### 4. PROBLEMS WITH BALANCE AND COORDINATION:

Dizziness, slow clumsy movements, inability to walk a straight line or balance on one foot with eyes closed.

**IF NO MEDICAL PERSONNEL ARE ON HAND AND AN INJURED ATHLETE HAS ANY OF THE ABOVE SYMPTOMS, HE OR SHE SHOULD BE SENT FOR APPROPRIATE MEDICAL CARE.**

## CHECKING FOR CONCUSSION

The presence of any of the signs or symptoms that are listed in this brochure suggest a concussion has most likely occurred. In addition to observation and direct questioning for symptoms, medical professionals have a number of other instruments to evaluate attention, processing speed, memory, balance, reaction time, and ability to think and analyze information (called executive brain function). These are the brain functions that are most likely to be adversely affected by a concussion and most likely to persist during the post concussion period.

---

*If an athlete seems "clear" he or she should be exercised enough to increase the heart rate and then evaluate if any symptoms return before allowing that athlete to practice or play.*

---

Computerized tests that can evaluate brain function are now being used by some medical professionals at all levels of sports from youth to professional and elite teams. They provide an additional tool to assist physicians in determining when a concussed athlete appears to have healed enough to return to school and play. This is especially helpful when dealing with those athletes denying symptoms in order to play sooner.

For non-medical personnel, the Centers for Disease Control and Prevention (CDC) has also developed a tool kit ("Heads Up: Concussion in High School Sports"), which has been made available to all high schools, and has information for coaches, athletes and parents. The NFHS is proud to be a co-sponsor of this initiative.

## PREVENTION

Although all concussions cannot be prevented, many can be minimized or avoided. Proper coaching techniques, good officiating of the existing rules, and use of properly fitted equipment can minimize the risk of head injury. Although the NFHS advocates the use of mouthguards in nearly all sports and mandates them in some, there is no convincing scientific data that their use will prevent concussions.

*Prepared by NFHS Sports Medicine Advisory Committee. 2009*

### References:

NFHS. Concussions. 2008 NFHS Sports Medicine Handbook (Third Edition). 2008: 77-82.  
NFHS. <http://www.nfhs.org>.

## National Federation of State High School Associations

PO Box 690 | Indianapolis, Indiana 46206  
Phone: 317-972-6900 | Fax: 317.822.5700  
[www.nfhs.org](http://www.nfhs.org)

National Federation of State  
High School Associations



# SUGGESTED GUIDELINES FOR MANAGEMENT OF CONCUSSION IN SPORTS

**EVEN SEEMINGLY MINOR CONCUSSIONS  
CAN HAVE DEVASTATING RESULTS**

## INTRODUCTION

Concussions are a common problem in sports and have the potential for serious complications if not managed correctly. Even what appears to be a "minor ding or bell ringer" has the real risk of catastrophic results when an athlete is returned to action too soon. The medical literature and lay press are reporting instances of death from "second impact syndrome" when a second concussion occurs before the brain has recovered from the first one regardless of how mild both injuries may seem.

At many athletic contests across the country, trained and knowledgeable individuals are not available to make the decision to return concussed athletes to play. Frequently, there is undo pressure from various sources (parents, player and coach) to return a valuable athlete to action. In addition, often there is unwillingness by the athlete to report headaches and other findings because the individual knows it would prevent his or her return to play.

Outlined below are some guidelines that may be helpful for parents, coaches and others dealing with possible concussions. Please bear in mind that these are general guidelines and must not be used in place of the central role that physicians and athletic trainers must play in protecting the health and safety of student-athletes.

## SIDELINE MANAGEMENT OF CONCUSSION

- 1. Did a concussion take place?** Based on mechanism of injury, observation, history and unusual behavior and reactions of the athlete, even without loss of consciousness, assume a concussion has occurred if the head was hit and even the mildest of symptoms occur. *(See other side for signs and symptoms)*
- 2. Does the athlete need immediate referral for emergency care?** If confusion, unusual behavior or responsiveness, deteriorating condition, loss of consciousness, or concern about neck and spine injury exist, the athlete should be referred at once for emergency care.
- 3. If no emergency is apparent, how should the athlete be monitored?** Every 5- 10 minutes, mental status, attention, balance, behavior, speech and memory should be examined until stable over a few hours. If appropriate medical care is not available, an athlete even with mild symptoms should be sent for medical evaluation.
- 4. No athlete suspected of having a concussion should return to the same practice or contest, even if symptoms clear in 15 minutes.**

## MANAGEMENT OF CONCUSSIONS AND RETURN TO PLAY

*(See "SIDELINE DECISION-MAKING" Below)*

Increasing evidence is suggesting that initial signs and symptoms, including loss of consciousness and amnesia, may not be very predictive of the true severity of the injury and the prognosis or outcome. More importance is being assigned to the duration of such symptoms and this, along with data showing symptoms may worsen some time after the head injury, has shifted focus to continued monitoring of the athlete. This is one reason why these guidelines no longer include an option to return an athlete to play even if clear in 15 minutes and why there is no discussion about the "Grade" of the concussion.

Any athlete who is removed from play because of a concussion should have medical clearance from an appropriate health care professional before being allowed to return to play or practice. The Second International Conference on Concussion held in Prague recommends an athlete should not return to practice or competition in sport until he or she is asymptomatic including after exercise.

Recent information suggests that mental exertion, as well as physical exertion, should be avoided until concussion symptoms have cleared. Premature mental or physical exertion may lead to more severe and more prolonged post concussion period. Therefore, the athlete should not study, play video games, do computer work or phone texting until his or her symptoms are resolving. Once symptoms are clear, the student-athlete should try reading for short peri-

ods of time. When 1-2 hours of studying can be done without symptoms developing, the athlete may return to school for short periods gradually increasing until a full day of school is tolerated without return of symptoms.

Once the athlete is able to complete a full day of school work, without PE or other exertion, the athlete can begin the gradual return to play protocol as outlined below. Each step increases the intensity and duration of the physical exertion until all skills required by the specific sport can be accomplished without symptoms. These recommendations have been based on the awareness of the increased vulnerability of the brain to concussions occurring close together and of the cumulative effects of multiple concussions on long-term brain function. Research is now revealing some fairly objective and relatively easy-to-use tests which appear to identify subtle residual deficits that may not be obvious from the traditional evaluation. These identifiable abnormalities frequently persist after the obvious signs of concussion are gone and appear to have relevance to whether an athlete can return to play in relative safety. The significance of these deficits is still under study and the evaluation instruments represent a work in progress. They may be helpful to the professional determining return to play in conjunction with consideration of the severity and nature of the injury; the interval since the last head injury; the duration of symptoms before clearing; and the level of play.

### SIDELINE DECISION-MAKING

1. No athlete should return to play (RTP) on the same day of concussion.
2. Any athlete removed from play because of a concussion must have medical clearance from an appropriate health care professional before he or she can resume practice or competition.
3. Close observation of athlete should continue for a few hours.
4. After medical clearance, RTP should follow a step-wise protocol with provisions for delayed RTP based on return of any signs or symptoms.

### A. ATHLETE MUST REMAIN ASYMPTOMATIC TO PROGRESS TO THE NEXT LEVEL.

### B. IF SYMPTOMS RECUR, ATHLETE MUST RETURN TO PREVIOUS LEVEL.

### C. MEDICAL CHECK SHOULD OCCUR BEFORE CONTACT.

### MEDICAL CLEARANCE RTP PROTOCOL

1. No exertional activity until asymptomatic.
2. When the athlete appears clear, begin low-impact activity such as walking, stationary bike, etc.
3. Initiate aerobic activity fundamental to specific sport such as skating or running, and may also begin progressive strength training activities.
4. Begin non-contact skill drills specific to sport such as dribbling, fielding, batting, etc.
5. Full contact in practice setting.
6. If athlete remains asymptomatic, he or she may return to game/play.